



FDA Approval of BELBUCA™ (CIII) (Buprenorphine HCl) Buccal Film for Chronic Pain Triggers Milestone Payment of \$50 Million to BioDelivery Sciences from Partner Endo Pharmaceuticals

October 26, 2015

Milestone Payment Provides BDSI Anticipated Cash Runway to Approximately Mid-2017

RALEIGH, N.C., Oct. 26, 2015 /PRNewswire/ -- BioDelivery Sciences International, Inc. (NASDAQ: BDSI) and its partner Endo Pharmaceuticals Inc. (Endo), a subsidiary of Endo International plc (NASDAQ: ENDP) (TSX: ENL), announced earlier today that the U.S. Food and Drug Administration (FDA) approved BELBUCA™ (CIII) (buprenorphine HCl) buccal film for use in patients with chronic pain severe enough to require daily, around-the-clock, long-term opioid treatment for which alternative treatment options are inadequate. FDA approval of BELBUCA has triggered a milestone payment to BDSI from Endo of \$50 million pursuant to the 2012 worldwide license and development agreement between BDSI and Endo for the development and commercialization of BELBUCA.

It is anticipated that BELBUCA may become commercially available in the U.S. during the first quarter of 2016, which may trigger additional milestone payments from Endo in the future if certain sales milestones are met. BDSI may also be entitled to receive tiered royalties on net sales of BELBUCA that start in the mid-teens.

Based on current operations and forecasts, BDSI estimates that the \$50 million dollar milestone payment from Endo, when combined with current cash on hand, provides BDSI with sufficient capital to operate its business to approximately the middle of 2017.

"The FDA approval of BELBUCA is a major milestone for BDSI and a tribute to the exceptional working relationship between the teams at Endo and BDSI. I want to thank all of our employees at BDSI for achieving our third product approval," said Dr. Mark A. Sirgo, President and Chief Executive Officer of BDSI.

"BELBUCA will provide a novel, alternative treatment option for millions of people suffering with chronic pain. Clinical trials demonstrated that twice daily administration of BELBUCA was well tolerated and controlled moderate to severe chronic pain. BELBUCA will also provide healthcare practitioners and their patients with a new buprenorphine treatment option. Buprenorphine is a Schedule III controlled substance, meaning that it has been defined as having lower abuse potential than Schedule II drugs, a category that includes most opioid analgesics," stated Dr. Andrew Finn, Executive Vice President of Product Development at BDSI.

Dr. Niraj Vasisht, Senior Vice President of Product Development and Chief Technical Officer at BDSI added, "Our patented BEMA delivery technology has enabled efficient and effective delivery of buprenorphine across the buccal mucosa and the seven approved dosages of BELBUCA will allow physicians to individualize titration and treatment based on the patient's analgesic needs."

INDICATION

BELBUCA™ (buprenorphine) buccal film is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioid formulations, reserve BELBUCA™ for use in patients for whom alternative treatment options (eg, non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- BELBUCA™ is not indicated as an as-needed (prn) analgesic.

IMPORTANT SAFETY INFORMATION about BELBUCA™

WARNING: ADDICTION, ABUSE, and MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL EXPOSURE; and NEONATAL OPIOID WITHDRAWAL SYNDROME

Addiction, Abuse, and Misuse

BELBUCA™ exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing BELBUCA™, and monitor patients regularly for the development of these behaviors or conditions.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of BELBUCA™. Monitor for respiratory depression, especially during initiation of BELBUCA™ or following a dose increase. Misuse or abuse of BELBUCA™ by chewing, swallowing, snorting, or injecting buprenorphine extracted from the buccal film will result in the uncontrolled delivery of buprenorphine and pose a significant risk of overdose and death.

Accidental Exposure

Accidental exposure to even one dose of BELBUCA™, especially by children, can result in a fatal overdose of buprenorphine.

Neonatal Opioid Withdrawal Syndrome

Prolonged use of BELBUCA™ during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available.

CONTRAINDICATIONS

BELBUCA™ is contraindicated in patients with:

- Significant respiratory depression
- Acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment
- Known or suspected gastrointestinal obstruction, including paralytic ileus
- Hypersensitivity (eg, anaphylaxis) to buprenorphine

WARNINGS AND PRECAUTIONS

Addiction, Abuse, and Misuse

- BELBUCA™ contains buprenorphine, a Schedule III controlled substance. As an opioid, BELBUCA™ exposes users to the risks of addiction, abuse, and misuse.
- Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing BELBUCA™, and monitor all patients receiving BELBUCA™ for the development of these behaviors or conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (eg, major depression). The potential for these risks should not, however, prevent the proper management of pain in any given patient. Patients at increased risk may be prescribed BELBUCA™, but use in such patients necessitates intensive counseling about the risks and proper use of BELBUCA™, along with intensive monitoring for signs of addiction, abuse, or misuse.
- Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed BELBUCA™ and in those who obtain the drug illicitly. Addiction can occur at recommended doses and if the drug is misused or abused.
- Abuse or misuse of BELBUCA™ by swallowing may cause choking, overdose, and death.
- Opioid agonists such as BELBUCA™ are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Strategies to reduce the risk include prescribing the drug in the smallest appropriate quantity and advising the patient on the proper disposal of unused drug.
- Contact a local state professional licensing board or state controlled substances authority for information on how to prevent and detect abuse or diversion of this product.
- BELBUCA™ should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain. Education and training programs that meet FDA requirements are offered by accredited CME/CE providers, and are available to prescribers of extended-release opioids at no or nominal cost. A list of these programs can be found at www.er-la-opioidREMS.com or by calling 1-800-503-0784. An FDA-approved patient counseling document for healthcare providers containing important safety information regarding the safe use, storage, and disposal of extended-release opioids can also be obtained or downloaded at www.er-la-opioidREMS.com.

Life-Threatening Respiratory Depression

- Serious, life-threatening, or fatal respiratory depression has been reported with the use of buprenorphine, even when used as recommended. Respiratory depression, from opioid use, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, supportive measures, and use of opioid antagonists, depending on the patient's clinical status.
- While serious, life-threatening or fatal respiratory depression can occur at any time during the use of BELBUCA™, the risk is greatest during initiation of therapy or following a dose increase. Closely monitor patients for respiratory depression when initiating therapy with BELBUCA™ and following dose increases.
- To reduce the risk of respiratory depression, proper dosing and titration of BELBUCA™ are essential. Overestimating the dose of BELBUCA™ when converting patients from another opioid product may result in fatal overdose with the first dose.
- Accidental exposure to BELBUCA™, especially in children, can result in respiratory depression and death due to an overdose of buprenorphine.

Neonatal Opioid Withdrawal Syndrome

Prolonged use of BELBUCA™ during pregnancy can result in withdrawal signs in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life threatening if not recognized and treated and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available.

- Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high-pitched cry,

tremor, vomiting, diarrhea, failure to gain weight; and there have been reports of convulsions, apnea, respiratory depression, and bradycardia. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn.

Risks due to Interactions with Central Nervous System Depressants

- Hypotension, profound sedation, coma, or respiratory depression may result if BELBUCA™ is used concomitantly with alcohol or other CNS depressants (eg, sedatives, anxiolytics, hypnotics, neuroleptics, other opioids).
- When considering the use of BELBUCA™ in a patient taking a CNS depressant, assess the duration of use of the CNS depressant and the patient's response, including the degree of tolerance that has developed to CNS depression. Additionally, evaluate the patient's use of alcohol or illicit drugs that cause CNS depression. If the decision to begin therapy with BELBUCA™ is made, start with a lower dosage of BELBUCA™, monitor patients for signs of sedation, respiratory depression, and hypotension, and consider using a lower dosage of the concomitant CNS depressant.

Risk of Life-Threatening Respiratory Depression in Elderly, Cachectic, and Debilitated Patients

- Monitor elderly, cachectic, or debilitated patients closely, particularly when initiating and titrating BELBUCA™ and when BELBUCA™ is given concomitantly with other drugs that depress respiration. Life-threatening respiratory depression is more likely to occur in these patients, as they may have altered pharmacokinetics or altered clearance compared with younger, healthier patients.

Risk of Apnea in Patients with Chronic Pulmonary Disease

- BELBUCA™-treated patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with substantially decreased respiratory reserve, hypoxia, hypercapnia, or preexisting respiratory depression are at increased risk of decreased respiratory drive, including apnea, even at recommended dosages of BELBUCA™. Therefore, closely monitor these patients, especially when initiating and titrating BELBUCA™. Alternatively, consider the use of alternative non-opioid analgesics in these patients.

QTc Prolongation

- BELBUCA™ has been observed to prolong the QTc interval in some subjects participating in clinical trials. Consider these observations in clinical decisions when prescribing BELBUCA™ to patients with hypokalemia, hypomagnesemia, or clinically unstable cardiac disease, including unstable atrial fibrillation, symptomatic bradycardia, unstable congestive heart failure, or active myocardial ischemia. Periodic electrocardiographic (ECG) monitoring is recommended in these patients. Avoid the use of BELBUCA™ in patients with a history of Long QT Syndrome or an immediate family member with this condition or those taking Class IA antiarrhythmic medications (e.g., quinidine, procainamide, disopyramide) or Class III antiarrhythmic medications (e.g., sotalolol, amiodarone, dofetilide), or other medications that prolong the QT interval.

Severe Hypotension

- BELBUCA™ may cause severe hypotension including orthostatic hypotension and syncope, in ambulatory patients. There is an increased risk in patients whose ability to maintain blood pressure has already been compromised by a reduced blood volume or concurrent administration of certain CNS depressant drugs (eg, phenothiazines or general anesthetics). Monitor these patients for signs of hypotension after initiating or titrating the dose of BELBUCA™. Avoid the use of BELBUCA™ in patients with circulatory shock, because BELBUCA™ may cause vasodilation that can further reduce cardiac output and blood pressure.

Risks of Use in Patients with Head Injury or Increased Intracranial Pressure

- In patients who may be susceptible to the intracranial effects of CO₂ retention (e.g., those with evidence of increased intracranial pressure or brain tumors), BELBUCA™ may reduce respiratory drive, and the resultant CO₂ retention can further increase intracranial pressure. Monitor such patients for signs of sedation and respiratory depression, particularly when initiating therapy with BELBUCA™.
- Opioids may also obscure the clinical course in a patient with a head injury. Avoid the use of BELBUCA™ in patients with impaired consciousness or coma.

Hepatotoxicity

- Cases of cytolytic hepatitis and hepatitis with jaundice have been observed in individuals receiving sublingual formulations of buprenorphine for the treatment of opioid dependence, both in clinical trials and in post-marketing adverse events reports. For patients at increased risk of hepatotoxicity (e.g., patients with a history of excessive alcohol intake, intravenous drug abuse or liver disease), obtain baseline liver enzyme levels and monitor periodically during treatment with

BELBUCA™.

Risk of Overdose in Patients With Moderate or Severe Hepatic Impairment

- In a pharmacokinetic study of subjects dosed with buprenorphine sublingual tablets, buprenorphine plasma levels were found to be higher and the half-life was found to be longer in subjects with moderate and severe hepatic impairment but not in subjects with mild hepatic impairment. For patients with severe hepatic impairment, a dose adjustment is recommended, and patients with moderate or severe hepatic impairment should be monitored for signs and symptoms of toxicity or overdose caused by increased levels of buprenorphine.

Anaphylactic/Allergic Reactions

- Cases of acute and chronic hypersensitivity to buprenorphine have been reported both in clinical trials and in post-marketing experience. The most common signs and symptoms include rashes, hives, and pruritus. Cases of bronchospasm, angioneurotic edema, and anaphylactic shock have been reported.

Risk of Use in Patients with Gastrointestinal Conditions

- BELBUCA™ may cause spasm of the sphincter of Oddi. Opioids may cause increases in the serum amylase. Monitor patients with biliary tract disease, including acute pancreatitis, for worsening symptoms.

Increased Risk of Seizures in Patients with Seizure Disorders

- Buprenorphine may increase the frequency of seizures in patients with seizure disorders and may increase the risk of seizures occurring in other clinical settings associated with seizures. Monitor patients with a history of seizure disorders for worsened seizure control during BELBUCA™ therapy.

Risks of Use in Cancer Patients with Oral Mucositis

- Cancer patients with oral mucositis may absorb buprenorphine more rapidly than intended and are likely to experience transiently higher plasma levels of the opioid. A dose reduction is recommended in these patients. Monitor carefully for signs and symptoms of toxicity or overdose caused by increased levels of buprenorphine.

Risks of Driving and Operating Machinery

- BELBUCA™ may impair the mental and physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery unless they are tolerant to side effects of BELBUCA™ and know how they will react to the medication.

ADVERSE REACTIONS

The most common adverse reactions (≥5%) reported by patients treated with BELBUCA™ in the clinical trials were nausea, constipation, headache, vomiting, fatigue, dizziness, somnolence, diarrhea, dry mouth, and upper respiratory tract infection.

About BioDelivery Sciences International

BioDelivery Sciences International, Inc. (NASDAQ:BDSI) is a specialty pharmaceutical company with a focus in the areas of pain management and addiction medicine. BDSI is utilizing its novel and proprietary BioErodible MucoAdhesive (BEMA®) technology and other drug delivery technologies to develop and commercialize, either on its own or in partnership with third parties, new applications of proven therapies aimed at addressing important unmet medical needs.

BDSI's development strategy focuses on utilization of the FDA's 505(b)(2) approval process. This regulatory pathway creates the potential for more timely and efficient approval of new formulations of previously approved therapeutics.

BDSI's particular area of focus is the development and commercialization of products in the areas of pain management and addiction. These are areas where BDSI believes its drug delivery technologies and products can best be applied to address critical unmet medical needs. BDSI's marketed products and those in development address serious and debilitating conditions such as breakthrough cancer pain, chronic pain, painful diabetic neuropathy and opioid dependence. BDSI's headquarters is located in Raleigh, North Carolina.

For more information, please visit or follow us:

Internet: www.bdsi.com

Cautionary Note on Forward-Looking Statements

This press release and any statements of employees, representatives and partners of BioDelivery Sciences International, Inc. (the "Company") related thereto contain, or may contain, among other things, certain "forward-looking statements" within the meaning of the Private Securities Litigation Reform Act of 1995. Such forward-looking statements involve significant risks and uncertainties. Such statements may include, without limitation, statements with respect to the Company's plans, objectives, projections, expectations and intentions and other statements identified by words such as "projects," "may," "will," "could," "would," "should," "believes," "expects," "anticipates," "estimates," "intends," "plans," "potential" or similar

expressions. These statements are based upon the current beliefs and expectations of the Company's management and are subject to significant risks and uncertainties, including those detailed in the Company's filings with the Securities and Exchange Commission. Actual results (including, without limitation, the results of (i) the anticipated commercial launch of BELBUCA; (ii) the potential for additional milestone payments and royalties to the Company from Endo and (iii) the timing for the use of the Company's current cash resources as described herein) may differ significantly from those set forth in the forward-looking statements. These forward-looking statements involve certain risks and uncertainties that are subject to change based on various factors (many of which are beyond the Company's control). The Company undertakes no obligation to publicly update any forward-looking statements, whether as a result of new information, future presentations or otherwise, except as required by applicable law.

BDSI[®], BEMA[®] and BUNAVAIL[®] are registered trademarks of BioDelivery Sciences International, Inc. The BioDelivery Sciences and BUNAVAIL logos are trademarks owned by BioDelivery Sciences International, Inc. BELBUCA[™] is owned by Endo International plc or one of its affiliates. All other trademarks and tradenames are owned by their respective owners.

© 2015 BioDelivery Sciences International, Inc. All rights reserved.

Logo - <http://photos.prnewswire.com/prnh/20110217/CL49801LOGO>

SOURCE BioDelivery Sciences International, Inc.

For further information: Investors: Matthew P. Duffy, Managing Director, LifeSci Advisors, LLC, 212-915-0685, matthew@lifesciadvisors.com; Al Medwar, Vice President, Marketing and Corporate Development, BioDelivery Sciences International, Inc., 919-582-9050, amedwar@bdsi.com; Media: Susan Forman/Laura Radocaj, Dian Griesel Int'l., 212-825-3210, sforman@dgicomm.com, lradocaj@dgicomm.com